



PATIENT DETAILS

Name
Address

Medicare No.

D.O.B.
Phone No.

INVESTIGATIONS REQUIRED

mpMRI Prostate Scan

CLINICAL NOTES

PSA = _____
PSA Free/Total Ratio = _____

REFERRING DOCTOR DETAILS

Doctor
Address

Provider
No.
Phone No.

COPY OF REPORT TO

REFERRING DOCTOR SIGNATURE

(Doctor's Name, Provider Number and Date are a legal requirement)

REFERRAL DATE

Additional Clinical Information

63541 mpMRI Prostate Scan for the diagnosis of prostate cancer

Must be suspected of developing prostate cancer based on:

- a digital rectal examination (DRE); or
- for a person 70 years or older, at least two PSA tests performed within an interval of 1-3 months have a PSA concentration of greater than 5.5 ng/L and the free/total PSA ratio is less than 25%; or
- for a person under 70 years, at least two prostate specific antigen (PSA) tests performed within an interval of 1-3 months have PSA concentration of greater than 3.0 ng/ml, and the free/total PSA ratio is less than 25%, or the repeat PSA exceeds 5.5 ng/L; or
- for a person under 70 years with a relevant family history, at least two PSA tests performed within an interval of 1-3 months have a PSA concentration greater than 2.0 ng/ml, and the free/total PSA ratio is less than 25%, or the repeat PSA exceeds 5.5 ng/L.

Note: Relevant family history is a first degree relative with prostate cancer or suspected of carrying a BRCA 1, BRCA 2 mutation.

Payable once per patient in a twelve month period.

63543 mpMRI Prostate Scan for the assessment of cancer

- the patient is under active surveillance following a confirmed diagnosis of prostate cancer by biopsy histopathology; and
- the patient is not planning or undergoing treatment for prostate cancer.

Claiming Notes:

- at least 12 months must have elapsed following confirmed diagnosis and then every third year thereafter
- at any time there is a clinical concern
- not to be used for the purpose of treatment, planning or for monitoring after treatment.

For patients who may require intravenous contrast and the patient is over 70 years old, or has suspected or known renal impairment please supply the serum levels within previous: 3 months for stable out-patient | 7 days for stable in-patient.

" eGFR= _____ Date of test: _____

Please bring this referral form at time of examination

Thank you for choosing South Coast Radiology

For Appointments: 1300 197 297 bookings@scr.com.au scr.com.au

① How to Book?



Booking your appointment is quick and easy.

1. Scan the QR code or,
2. Visit scr.com.au to book online

OR

1300 197 297

Call one of our friendly team members to schedule your appointment.

② Your Appointment

Date: _____

Time: _____

Clinic location: _____

Please arrive 15 minutes prior to your appointment time.

You will receive detailed preparation instructions for your appointment at the time of booking.

③ Accessing your Images

South Coast Radiology offers iVue – our free Patient App and secure digital portal giving you convenient access to your medical imaging performed at South Coast Radiology. View your images anytime from your smartphone, tablet, or web browser. Share them with your healthcare providers or family, and easily request appointments when needed. Download our free iVue Patient App via the App store on your smart phone.

Please Note: Your images will reach your account 7 days after your appointment.

④ Clinic Locations

South Coast Radiology proudly operates clinics across the Gold & Tweed Coast – including our dedicated Women's Imaging Centre and two regional locations.



To find the clinic most convenient to you, scan the QR code to view our full list of locations.

Important Medicare Advice: Your referrer has recommended you use South Coast Radiology. You may choose another provider, but please discuss this with your referrer first.

Scan Preparation:

To achieve a successful MRI Prostate scan, it is important that your bowel is empty for the procedure.

You will be given 2 Dulcolax tablets to take the night before the procedure. These tablets should cause evacuation of your bowel in the morning of the day of your procedure. Please inform our staff prior to the procedure if evacuation of your bowel did not occur.

We ask that you do not have an ejaculation for 48 hours prior to appointment.

You must not have anything to eat, drink, smoke, or chew for 6 hours prior to appointment.

Do you have any of the following?

- Metal in your eye
- Cardiac pacemaker
- Epicardial wires
- Brain aneurysm clip
- I.V.C. filter
- Inner ear implant
- Claustrophobia
- Neurostimulator
- Brain shunt
- Continuous glucose monitoring devices
- Are you currently pregnant?

What to Expect:

Before Your Scan:

An intramuscular injection of buscopan or glucagon will be administered approximately 30 minutes prior to your MRI to relax the bowel and minimise motion during scan.

You may be asked to change into a patient gown prior to your scan. You will need to remove all jewellery; your valuables will be stored securely during your scan.

Our MRI radiographer will explain the examination in detail and check your MRI Safety Questionnaire. Once this has been checked by the MRI radiographers that you are safe to have an MRI scan, they will prepare you for the scan by inserting an IV cannula for the MRI contrast to be injected during this test.

A small plastic tube (IV cannula) will be inserted into a vein in your arm so we can administer contrast during the scan. The contrast helps enhance the MRI images and provides our radiologist with more detailed information.

During Your Scan:

During the MRI scan, you will be positioned on your back on the MRI bed and your pelvis and abdomen will be 'inside' the MRI scanner. It's essential that you remain perfectly still throughout the scan. Movement can impact image quality. If you require pain medication to comfortably lie on your back, please take it before arriving.

The MRI makes loud noises, but you'll be given protective headphones. We can also play music of your choice - just let our staff know. Some patients may feel claustrophobic during the scan. If this is a concern, please let us know at the time of booking so we can assist accordingly. While you'll be alone in the room, our technician will be monitoring you the entire time. You'll be able to communicate via intercom and will be given a buzzer in case you need assistance.

After the initial images are taken, the contrast will be injected through your IV cannula. You may feel a cool sensation or slight nausea briefly. More scans will be taken following the injection.

The full scan usually takes 20–30 minutes.

After Your Scan:

You can resume to your usual daily activities immediately. Your images will be reviewed by a specialist Radiologist, and your referring doctor will receive a detailed report. You can also view your results and images through our secure online patient portal,

which you can register for during your visit.



**SOUTH COAST
RADIOLOGY**

Trusted by Doctors, Preferred by Patients